

COMPARISON OF POISSON STRUCTURES AND POISSON-LIE DYNAMICAL r -MATRICES

BENJAMIN ENRIQUEZ, PAVEL ETINGOF, AND IAN MARSHALL

ABSTRACT. We construct a Poisson isomorphism between the formal Poisson manifolds $\mathfrak{g}^*$ and G^* , where $\mathfrak{g}$ is a finite dimensional quasitriangular Lie bialgebra. Here $\mathfrak{g}^*$ is equipped with its Lie-Poisson (or Kostant-Kirillov-Souriau) structure, and G^* with its Poisson-Lie structure. We also quantize the Poisson-Lie dynamical r -matrices of Balog-Fehér-Palla.

INTRODUCTION AND MAIN RESULTS

We construct Poisson isomorphisms between the formal Poisson manifolds $\mathfrak{g}^*$ and G^* , where $\mathfrak{g}$ is a finite dimensional quasitriangular Lie bialgebra. Here $\mathfrak{g}^*$ is equipped with its Lie-Poisson (or Kostant-Kirillov-Souriau) structure, and G^* with its Poisson-Lie structure.

This result may be viewed as a generalization of the formal version of [GW] (later reproved in [Al], Theorem 1, and [Bo]), where Ginzburg and Weinstein construct a Poisson diffeomorphism between the real Poisson manifolds $\mathfrak{k}^*$ and K^* , where K is a compact Lie group and $\mathfrak{k}$ is its Lie algebra. It can also be viewed as a new result in the subject of linearization of Poisson structures; e.g., in contrast with our result, it has been shown in [Ch] that not all Poisson structures on Poisson-Lie groups are linearizable.

Our result relies on constructing a map $g(\lambda) : \mathfrak{g}^* \rightarrow G$ satisfying a differential equation. We give two constructions of $g(\lambda)$: the first one uses the theory of the classical Yang-Baxter equation and gauge transformations; the second one relies on the theory of quantization of Lie bialgebras.

We then apply this result to the quantization of Poisson-Lie dynamical r -matrices introduced in [FM1].

We now describe our results in more detail.

0.1. Comparison of Poisson structures. Let $(\mathfrak{g}, r)$ be a finite dimensional quasitriangular Lie bialgebra over a field $\mathbf{k}$ of characteristic 0. Recall that this means that $\mathfrak{g}$ is a Lie algebra, $r \in \mathfrak{g}^{\otimes 2}$, $t := r + r^{2,1}$ is invariant, and $\text{CYB}(r) := [r^{1,2}, r^{1,3}] + [r^{1,2}, r^{2,3}] + [r^{1,3}, r^{2,3}] = 0$. The Lie cobracket on $\mathfrak{g}$ is defined by $\delta(x) := [x \otimes 1 + 1 \otimes x, r]$ for any $x \in \mathfrak{g}$.

The Lie bracket on $\mathfrak{g}$ defines a linear Poisson structure on $\mathfrak{g}^*$; we also denote by $\mathfrak{g}^*$ the formal neighborhood of the origin in this vector space, which is then a formal Poisson manifold. On the other hand, we denote by G^* the formal Poisson-Lie group with Lie bialgebra $\mathfrak{g}^*$. Our main result is:

Theorem 0.1. *The formal Poisson manifolds $\mathfrak{g}^*$ and G^* are isomorphic.*

We now explicitly describe an isomorphism $\mathfrak{g}^* \rightarrow G^*$. Denote by G the formal group with Lie algebra $\mathfrak{g}$ and by $\text{Map}_0(\mathfrak{g}^*, G)$ the space of formal maps $g : \mathfrak{g}^* \rightarrow G$, such that $g(0) = 1$; this is the space of maps of the form $e^{x(\lambda)}$, where $x(\lambda) \in \mathfrak{g} \otimes \widehat{S}(\mathfrak{g})_{>0}$.¹

¹We denote by $\widehat{S}(\mathfrak{g})$ the degree completion $\widehat{\oplus}_{k \geq 0} S^k(\mathfrak{g})$ of the symmetric algebra $S(\mathfrak{g})$, and set $\widehat{S}(\mathfrak{g})_{>i} = \widehat{\oplus}_{k > i} S^k(\mathfrak{g})$.

$\text{Map}_0(\mathfrak{g}^*, G)$ has a group structure, defined by $(g_1 * g_2)(\lambda) := g_2(\text{Ad}^*(g_1(\lambda))(\lambda))g_1(\lambda)$. Its subspace of all maps $g(\lambda)$ such that

$$g_1^{-1}d_2(g_1)(\lambda) - g_2^{-1}d_1(g_2)(\lambda) + \langle \text{id} \otimes \text{id} \otimes \lambda, [g_1^{-1}d_3(g_1)(\lambda), g_2^{-1}d_3(g_2)(\lambda)] \rangle = 0 \quad (1)$$

is a subgroup $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$. Here $g_1^{-1}d_2(g_1)(\lambda) = \sum_i g^{-1}\partial_{\varepsilon^i}g(\lambda) \otimes e_i$ is viewed as a formal function $\mathfrak{g}^* \rightarrow \mathfrak{g}^{\otimes 2}$, (ε^i) , (e_i) are dual bases of $\mathfrak{g}^*$ and $\mathfrak{g}$, $g_i^{-1}d_j(g_i) = (g_1^{-1}d_2(g_1))^{i,j}$ and $\partial_\xi g(\lambda) = (d/d\varepsilon)|_{\varepsilon=0}g(\lambda + \varepsilon\xi)$. We will also denote by $g_{13}^{-1}d_2(g_{13})$ the same quantity, viewed as an element of $\mathfrak{g}^{\otimes 2} \otimes \widehat{S}(\mathfrak{g})$.

We have a group morphism $\theta : \text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G) \rightarrow \text{Aut}_1(\mathfrak{g})^{\text{op}}$ to the group (with opposite structure) of Poisson automorphisms of $\mathfrak{g}^*$ with differential at 0 equal to the identity, taking $g(\lambda)$ to the automorphism $\lambda \mapsto \text{Ad}^*(g(\lambda))(\lambda)$.

Proposition 0.2. *There exists a formal map $g(\lambda) \in \text{Map}_0(\mathfrak{g}^*, G)$, such that*

$$(g_1)^{-1}d_2(g_1) - (g_2)^{-1}d_1(g_2) + \text{Ad}(g \otimes g)^{-1}(r_0) + \langle \text{id} \otimes \text{id} \otimes \lambda, [(g_1)^{-1}d_3(g_1), (g_2)^{-1}d_3(g_2)] \rangle = \rho_{\text{AM}}. \quad (2)$$

(identity of formal maps $\mathfrak{g}^* \rightarrow \wedge^2(\mathfrak{g})$).² Here $r_0 = (r - r^{2,1})/2$, and ρ_{AM} is the Alekseev-Meinrenken r -matrix ([AM, BDF]) given by

$$\rho_{\text{AM}}(\lambda) = (\text{id} \otimes \varphi(\text{ad } \lambda^\vee))(t),$$

where $\lambda^\vee = (\lambda \otimes \text{id})(t)$ and $\varphi(z) := -\frac{1}{z} + \frac{1}{2} \coth \frac{z}{2}$.

The group $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ acts simply and transitively on the space of solutions $g(\lambda)$ of (2), as follows: $(\alpha * g)(\lambda) = g(\text{Ad}^*(\alpha(\lambda))(\lambda))\alpha(\lambda)$.

Proposition 0.3. *Let $g(\lambda) \in \text{Map}_0(\mathfrak{g}^*, G)$ be as in Proposition 0.2. There exists a unique isomorphism $g^*(\lambda) : \mathfrak{g}^* \rightarrow G^*$, defined by the identity*

$$g(\lambda)e^{\lambda^\vee}g(\lambda)^{-1} = L(g^*(\lambda))R(g^*(\lambda))^{-1}.$$

Here $L, R : G^* \rightarrow G$ are the formal group morphisms corresponding to the Lie algebra morphisms $L, R : \mathfrak{g}^* \rightarrow \mathfrak{g}$ given by $L(\lambda) := (\lambda \otimes \text{id})(r)$, $R(\lambda) := -(\lambda \otimes \text{id})(r^{2,1})$.

In other words, we have (non-Poisson) formal manifold isomorphisms $\mathfrak{g}^* \xrightarrow{a} G \xleftarrow{b} G^*$, $a(\lambda) = g(\lambda)e^{\lambda^\vee}g(\lambda)^{-1}$, $b(g^*) = L(g^*)R(g^*)^{-1}$, and $g^*(\lambda) = b^{-1} \circ a(\lambda)$.

The isomorphism $(\alpha * g)^*(\lambda) : \mathfrak{g}^* \rightarrow G^*$ corresponding to $(\alpha * g)(\lambda)$ is such that $(\alpha * g)^*(\lambda) = g^*(\theta(\alpha)(\lambda))$.

It follows that the set of all isomorphisms $\mathfrak{g}^* \rightarrow G^*$ constructed in Propositions 0.2, 0.3 is a principal homogeneous space under the image of $\theta : \text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G) \rightarrow \text{Aut}_1(\mathfrak{g}^*)$. When $\mathfrak{g}$ is semisimple, any derivation $\mathfrak{g} \rightarrow S^k(\mathfrak{g})$ is inner, so θ is surjective. So in that case, our construction yields all the Poisson isomorphisms $\mathfrak{g}^* \rightarrow G^*$ taking 0 to 1 and with differential at 0 equal to the identity.

0.2. Quantization of Poisson-Lie dynamical r -matrices. In [BFP, FM1], there was introduced the Poisson-Lie dynamical r -matrix

$$\rho_{\text{FM}}(g^*) = \left(\text{id} \otimes \left(\nu \frac{\text{id} + a(g^*)^{2\nu}}{\text{id} - a(g^*)^{2\nu}} - \frac{1}{2} \frac{\text{id} + a(g^*)}{\text{id} - a(g^*)} \right) \right)(t).$$

Here $a(g^*) : G^* \rightarrow \text{GL}(\mathfrak{g})$ is defined by $a(g^*) = \text{Ad}(L(g^*)R(g^*)^{-1})$ and ν is a fixed element in $\mathbf{k}$. Then ρ_{FM} is a formal map $G^* \rightarrow \wedge^2(\mathfrak{g})$, i.e., an element of $\wedge^2(\mathfrak{g}) \otimes \mathcal{O}_{G^*}$. It is a Poisson-Lie classical dynamical r -matrix for $(G^*, \mathfrak{g}, Z_\nu)$, where $Z_\nu = (\nu^2 - \frac{1}{4})[t^{1,2}, t^{2,3}]$ (see also [EEM]).

²We view $\wedge^n(\mathfrak{g})$ as a subspace of $\mathfrak{g}^{\otimes n}$.

A quantization of this dynamical r -matrix is the data of a quantization $U_{\hbar}(\mathfrak{g})$ of the Lie bialgebra $\mathfrak{g}$, and a pair $(\bar{J}, \bar{\Phi})$, where:

- $\bar{\Phi}$ is an associator for $U_{\hbar}(\mathfrak{g})$, i.e., $\bar{\Phi} \in U_{\hbar}(\mathfrak{g})^{\otimes 3}$ satisfies^{3,4} the pentagon equation $\bar{\Phi}^{2,3,4} \bar{\Phi}^{1,23,4} \bar{\Phi}^{1,2,3} = \bar{\Phi}^{1,2,34} \bar{\Phi}^{12,3,4}$, the invariance condition $[\bar{\Phi}, (\Delta \otimes \text{id}) \circ \Delta(a)] = 0$ for any $a \in U_{\hbar}(\mathfrak{g})$, and the expansion $\bar{\Phi} = 1 + \hbar^2 \phi_2$, where $\phi_2 \in U_{\hbar}(\mathfrak{g})^{\otimes 3}$ satisfies $\sum_{\sigma \in \mathfrak{S}_3} \text{sign}(\sigma) \sigma((\phi_2)|_{\hbar=0}) = Z_{\nu}$;
- $\bar{J} \in U_{\hbar}(\mathfrak{g})^{\otimes 2} \hat{\otimes} U_{\hbar}(\mathfrak{g})'$ satisfies the dynamical twist equation $\bar{J}^{2,3,4} \bar{J}^{1,23,4} \bar{\Phi}^{1,2,3} = \bar{J}^{1,2,34} \bar{J}^{12,3,4}$, the invariance condition $[\bar{J}, (\Delta \otimes \text{id}) \circ \Delta(a)] = 0$ for any $a \in U_{\hbar}(\mathfrak{g})$, and the expansion $\bar{J} = 1 + \hbar \bar{j}_1$, where $\bar{j}_1 \in U_{\hbar}(\mathfrak{g})^{\otimes 2} \hat{\otimes} U_{\hbar}(\mathfrak{g})'$ satisfies $(\bar{j}_1 - \bar{j}_1^{2,1,3})|_{\hbar=0} = \rho_{\text{FM}}$.

Theorem 0.4. *Such a quantization exists (see Theorem 2.1).*

1. CONSTRUCTION OF ISOMORPHISMS $\mathfrak{g}^* \rightarrow G^*$

Theorem 0.1 follows from Propositions 0.2 and 0.3. In this Section, we give two proofs of these propositions: (a) a Poisson geometric proof (Section 1.1), and (b) a proof based on the theory of quantization of Lie bialgebras (Section 1.2).

1.1. Geometric construction.

1.1.1. *Construction of $g(\lambda)$ (proof of Proposition 0.2).* One checks that $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ is a prounipotent Lie group with Lie algebra $\{\alpha \in \mathfrak{g} \otimes \hat{S}(\mathfrak{g})_{\geq 1} \mid \text{Alt} \circ d(\alpha) = 0\}$. This Lie algebra is isomorphic to $(\hat{S}(\mathfrak{g})_{>1}, \{-, -\})$ under $d : f \mapsto d(f)$.⁵

Let us denote by $\mathcal{G}$ the set of all $g \in \text{Map}_0(\mathfrak{g}^*, G)$ satisfying (2). This is a subvariety of the proalgebraic variety $\text{Map}_0(\mathfrak{g}^*, G)$. One checks that $(\hat{S}(\mathfrak{g})_{>1}, \{-, -\})$ acts by vector fields on $\mathcal{G}$, by

$$g^{-1} \delta_f(g) = \langle \text{id} \otimes \text{id} \otimes \lambda, [d_3(f_2), g_{12}^{-1} d_3(g_{12})] \rangle - d_1(f_2) \in \mathfrak{g} \otimes \hat{S}(\mathfrak{g})_{>0}, \quad (3)$$

and that the right infinitesimal action of $\text{Map}_0(\mathfrak{g}^*, G)$ on itself is given by the same formula. It follows that the right action of $\text{Map}_0(\mathfrak{g}^*, G)$ on itself restricts to an action of $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ on $\mathcal{G}$.

We now prove that if $\mathcal{G}$ is nonempty, then $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ acts simply and transitively on $\mathcal{G}$.

Let us show that the action is simple. If $g, g' \in \mathcal{G}$ and $\alpha \in \text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ are such that $g * \alpha = g'$, then let $a := \log(\alpha)$. Assume that $a \neq 0$ and let n be the smallest integer such that the component a_n of a in $\mathfrak{g} \otimes S^n(\mathfrak{g})$ is $\neq 0$. Then $\log(g') - \log(g) = a_n$ modulo $\mathfrak{g} \otimes \hat{S}(\mathfrak{g})_{>n}$, which implies that $g \neq g'$.

Let us now prove that the action is transitive. Let $g, g' \in \mathcal{G}$. Set $A := \log(g)$, $A' := \log(g')$. Then $A, A' \in \mathfrak{g} \otimes \hat{S}(\mathfrak{g})_{>0}$. Assume that $A \neq A'$ and let n be the smallest integer such that the component of $A' - A$ in $\mathfrak{g} \otimes S^n(\mathfrak{g})$ is nonzero; we denote by $(A' - A)_n$ this component. Comparing equations (2) for g and g' , we get $\text{Alt} \circ d((A' - A)_n) = 0$. It follows that there exists $a \in S^{n+1}(\mathfrak{g})$, such that $(A' - A)_n = \text{Alt} \circ d(a)$, i.e., $(A' - A)_n \in \text{Lie Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$. Let $\exp_*$ be the exponential map of $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ and $\alpha := \exp_*((A' - A)_n) \in \text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$. Then $\log(\alpha * g) = A(\text{Ad}^*(\alpha(\lambda))(\lambda)) + (A' - A)_n$ modulo $\mathfrak{g} \otimes \hat{S}(\mathfrak{g})_{\geq n+1}$; therefore the difference of logarithms⁶ of $\alpha * g$ and g' coincide modulo $\mathfrak{g} \otimes \hat{S}(\mathfrak{g})_{\geq n+1}$. Working by successive approximations, we construct $\beta \in \text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ such that $\beta * g = g'$. This proves that the action is transitive.

³We denote by $\hat{\otimes}$ the topological tensor product, defined as follows: if V, W are topological vector spaces of the form $V = V_0[[x_1, \dots, x_n]]$, $W = W_0[[y_1, \dots, y_m]]$, then $V \hat{\otimes} W := V_0 \otimes W_0[[x_1, \dots, y_m]]$.

⁴We set $\bar{\Phi}^{12,3,4} = (\Delta \otimes \text{id} \otimes \text{id})(\bar{\Phi})$, etc.

⁵For $\rho = \alpha \otimes f \in \wedge^n(\mathfrak{g}) \otimes \hat{S}(\mathfrak{g})$, we set $d(\rho) := \sum_i \alpha \otimes e_i \otimes (d/d\varepsilon)|_{\varepsilon=0} f(\lambda + \varepsilon e^i)$ and if $\xi \in \wedge^{n-1}(\mathfrak{g}) \otimes \mathfrak{g}$, $\text{Alt}(\xi \otimes f) := (\xi + \xi^{2, \dots, n, 1} + \dots + \xi^{n, 1, \dots, n-1}) \otimes f$.

⁶Here the logarithm is the inverse of the ordinary exponential map $\mathfrak{g} \otimes \hat{S}(\mathfrak{g}) \rightarrow \text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$.

Let us now prove that $\mathcal{G}$ is nonempty. Recall that $r_0 = (r - r^{2,1})/2$. If $g \in \text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$, set $(r_0)^g := \text{l.h.s. of (2)}$.

Lemma 1.1. *Assume that $\log(g) = -r/2$ modulo $\mathfrak{g} \otimes \widehat{S}(\mathfrak{g})_{\geq 1}$. Set $\rho := (r_0)^g$ and assume that $\rho = \rho_{\text{inv}} + \alpha$, where $\rho_{\text{inv}}, \alpha \in \wedge^2(\mathfrak{g}) \otimes \widehat{S}(\mathfrak{g})$, ρ_{inv} is $\mathfrak{g}$ -invariant and $\alpha \in \mathfrak{g} \otimes \widehat{S}(\mathfrak{g})_{\geq n}$. Then*

$$\text{CYB}(\rho) - \text{Alt}(d\rho) = Z^{1,2,3} - \frac{1}{2}[r^{1,234}, \alpha^{2,3,4}]$$

modulo $\mathfrak{g}^{\otimes 3} \otimes \widehat{S}(\mathfrak{g})_{\geq n+1}$.

Proof of Lemma. This statement can be proved directly. It can also be viewed as the classical limit of the following statement. Let $U = U(\mathfrak{g})[[\hbar]]$, let $J \in (U^{\widehat{\otimes} 2})^\times$, and let $\Phi := (J^{1,2} J^{12,3})^{-1} J^{2,3} J^{1,23}$. If $K \in (U^{\widehat{\otimes} 2})^\times$ and we set $\bar{\Phi} := (K^{1,23})^{-1} (K^{2,3})^{-1} J^{1,2} K^{12,3}$, then $\bar{\Phi}$ satisfies

$$\bar{\Phi}^{2,3,4} \bar{\Phi}^{1,23,4} \bar{\Phi}^{1,2,3} = (\bar{\Phi}^{2,3,4}, (K^{1,234})^{-1}) \bar{\Phi}^{1,2,34} \bar{\Phi}^{12,3,4} (K^{123,4}, \Phi^{1,2,3}).$$

Here $(a, b) = aba^{-1}b^{-1}$. The statement of the lemma is recovered when $J, K \in (U^{\widehat{\otimes} 2})^\times$ have the form $1 - \hbar r/2 + o(\hbar)$, and K is admissible with classical limit $g(\lambda)$: the contribution of $(\bar{\Phi}^{2,3,4}, (K^{1,234})^{-1})$ is $-\frac{1}{2}[r^{1,234}, \alpha^{2,3,4}]$, while the commutator $(K^{123,4}, \Phi^{1,2,3})$ does not contribute (as the classical limit of Φ is proportional to Z and hence invariant). $\square$

Let us now prove that $\mathcal{G}$ is nonempty. We will construct a sequence $g_n \in \text{Map}_0(\mathfrak{g}^*, G)$, such that $\rho_n := (r_0)^{g_n}$ satisfies $\rho_n = \rho_{\text{AM}}$ modulo $\wedge^3(\mathfrak{g}) \otimes \widehat{S}(\mathfrak{g})_{\geq n+1}$.

If $n = 0$, we set $g_0 := \exp(-r/2)$, then $\rho_0 = 0 = \rho_{\text{AM}}$ modulo $\wedge^3(\mathfrak{g}) \otimes \widehat{S}(\mathfrak{g})_{\geq 1}$. Assume that we determined g_n and let us construct g_{n+1} . Set $\alpha := \rho_n - \rho_{\text{AM}}$, and let α_{n+1} be the component of α in $\wedge^2(\mathfrak{g}) \otimes S^{n+1}(\mathfrak{g})$. Then Lemma 1.1 implies that

$$\text{CYB}(\rho_{\text{AM}} + \alpha_{n+1}) - \text{Alt}(d(\rho_{\text{AM}} + \alpha_{n+1})) = Z^{1,2,3} - \frac{1}{2}[r^{1,234}, \alpha_{n+1}^{2,3,4}] \mod \wedge^3(\mathfrak{g}) \otimes S^{n+1}(\mathfrak{g}).$$

Since ρ_{AM} satisfies the modified classical dynamical Yang-Baxter equation, the component in $\wedge^3(\mathfrak{g}) \otimes S^n(\mathfrak{g})$ of this identity yields $\text{Alt}(d\alpha_{n+1}) = 0$, so that we have $\alpha_{n+1} = \text{Alt}(d\beta)$ for some $\beta \in \mathfrak{g} \otimes S^{n+2}(\mathfrak{g})$. Then we set $g_{n+1} := \exp_*(-\beta) * g_n$, $\rho_{n+1} := (r_0)^{g_{n+1}}$. Then $\rho_{n+1} - \rho_{\text{AM}} = \rho_n - \rho_{\text{AM}} - \text{Alt}(d\beta) = \alpha_{n+1} - \text{Alt}(d\beta) = 0$ modulo $\wedge^3(\mathfrak{g}) \otimes \widehat{S}(\mathfrak{g})_{\geq n+1}$. By successive approximations, we then construct g such that $(r_0)^g = \rho_{\text{AM}}$. So $\mathcal{G}$ is nonempty.

1.1.2. Poisson isomorphism $\mathfrak{g}^* \rightarrow G^*$ (proof of Proposition 0.3). In this section, we show that the fact that $g(\lambda)$ satisfies (2) implies that $\lambda \mapsto g^*(\lambda)$ is a Poisson isomorphism. We will freely use the formalism of differential geometry, even though we work in the formal setup; the computations below make sense (and prove the desired result) when working over an Artinian $\mathbf{k}$ -ring.

According to [STS], the image of the Poisson bracket $\{-, -\}_{G^*}$ on G^* under the formal isomorphism $b : G^* \rightarrow G$ is the Poisson bracket on G

$$\{F, H\}_G(g) = \langle (d_{\mathbf{R}} - d_{\mathbf{L}})F(g) \otimes d_{\mathbf{L}}H(g), r \rangle + \langle (d_{\mathbf{R}} - d_{\mathbf{L}})F(g) \otimes d_{\mathbf{R}}H(g), r^{2,1} \rangle,$$

where $g \in G$, F, H are functions on G , $d_{\mathbf{L}}F(g), d_{\mathbf{R}}F(g) \in \mathfrak{g}^*$ are the left and right differentials defined by $\langle d_{\mathbf{L}}F(g), a \rangle = (d/d\varepsilon)|_{\varepsilon=0} F(e^{\varepsilon a}g)$, $\langle d_{\mathbf{R}}F(g), a \rangle = (d/d\varepsilon)|_{\varepsilon=0} F(g e^{\varepsilon a})$ for any $a \in \mathfrak{g}$.

For $\xi \in \mathfrak{g}^*$, define $F_\xi \in \mathcal{O}_G$ by $F_\xi(g) = \langle \xi, \log(g) \rangle$.

Lemma 1.2. $(d_{\mathbf{L}}F_\xi)(g) = f(\frac{1}{2}\text{ad}^*(\log g))(\xi)$ and $(d_{\mathbf{R}}F_\xi)(g) = f(-\frac{1}{2}\text{ad}^*(\log g))(\xi)$, where $f(z) = ze^z/(\sinh z)$, and ad^* denotes the coadjoint action of $\mathfrak{g}$ on $\mathfrak{g}^*$.

Proof. Set $x := \log(g)$, let $a \in \mathfrak{g}$, and set $\tilde{a} := (d/d\varepsilon)|_{\varepsilon=0} \log(e^{\varepsilon a}e^x)$. The coefficient in ε of $e^{\varepsilon a}e^x = e^{x+\varepsilon\tilde{a}+O(\varepsilon^2)}$ yields $ae^x = \sum_{n \geq 1} (n!)^{-1} \sum_{k=0}^{n-1} x^k \tilde{a} x^{n-1-k}$. Applying $\text{ad}(x)$ to this relation, we get $\tilde{a} = f(-\frac{1}{2}\text{ad } x)(a)$. Now $\langle d_{\mathbf{L}}F_\xi(g), a \rangle = (d/d\varepsilon)|_{\varepsilon=0} F_\xi(e^{\varepsilon a}g) = \langle \xi, \tilde{a} \rangle =$

$\langle \xi, f(-\frac{1}{2} \text{ad } x)(a) \rangle = \langle f(\frac{1}{2} \text{ad}^* x)(\xi), a \rangle$, which yields the first formula. On the other hand, we have for any function F on G , $d_{\mathbf{R}} F(g) = \text{Ad}^*(g^{-1})(d_{\mathbf{L}} F(g))$, where Ad^* is the coadjoint action of G on $\mathfrak{g}^*$, hence $d_{\mathbf{R}} F_{\xi}(g) = \text{Ad}^*(e^{-x})f(\frac{1}{2} \text{ad}^* x)(\xi) = f(-\frac{1}{2} \text{ad}^* x)(\xi)$. $\square$

Then we get

$$\{F_{\xi}, F_{\eta}\}_G(g) = \langle \text{ad}^*(x)(\xi) \otimes \text{ad}^*(x)(\eta), (\text{id} \otimes \varphi(\text{ad}(x)))(t) + r_0 \rangle - \langle \text{ad}^*(x)(\xi) \otimes \eta, t \rangle, \quad (4)$$

where $x = \log(g)$ and φ and r_0 are as in Proposition 0.2.

On the other hand, let $\{-, -\}'_G$ be the image of the Poisson bracket $\{-, -\}_{\mathfrak{g}^*}$ by $a : \mathfrak{g}^* \rightarrow G$. Set $f_{\xi}(\lambda) := F_{\xi} \circ a(\lambda)$, then $f_{\xi}(\lambda) = \langle \xi \otimes \lambda, (\text{Ad}(g(\lambda)) \otimes \text{id})(t) \rangle$. If $f \in \mathcal{O}_{\mathfrak{g}^*}$, $\lambda \in \mathfrak{g}^*$, define $df(\lambda) \in \mathfrak{g}$ by $\langle \alpha, df(\lambda) \rangle = (d/d\varepsilon)|_{\varepsilon=0} f(\lambda + \varepsilon \alpha)$ for any $\alpha \in \mathfrak{g}^*$. We have $\{f, g\}_{\mathfrak{g}^*}(\lambda) = \langle \lambda, [df(\lambda), dg(\lambda)] \rangle$. Then $df_{\xi}(\lambda) = \langle \xi \otimes \text{id}, A(\lambda) \rangle$, where

$$A(\lambda) = (\text{Ad}(g(\lambda)) \otimes \text{id})(t) + [(d_2 g_1) g_1^{-1}, \text{Ad}(g(\lambda))(\lambda^{\vee}) \otimes 1].$$

So

$$\{f_{\xi}(\lambda), f_{\eta}(\lambda)\} = \langle \xi \otimes \eta \otimes \lambda, [A^{1,3}(\lambda), A^{2,3}(\lambda)] \rangle.$$

This decomposes as the sum of four terms (we set $\bar{x} = \text{Ad}(g(\lambda))(\lambda^{\vee})$):

- (a) $\langle \xi \otimes \eta \otimes \lambda, (\text{Ad}(g(\lambda))^{\otimes 2} \otimes \text{id})([t^{13}, t^{23}]) \rangle = -\langle \xi \otimes \eta, (\text{ad}(\bar{x}) \otimes \text{id})(t) \rangle$;
- (b) $\langle \xi \otimes \eta \otimes \lambda, [(d_3 g_1) g_1^{-1}, \text{Ad}(g_1(\lambda))(\lambda^{\vee})], \text{Ad}(g_2(\lambda))(t^{23}) \rangle = \langle \text{ad}^*(\bar{x})^{\otimes 2}(\xi \otimes \eta), -\text{Ad}(g)^{\otimes 2}(g_1^{-1} d_2(g_1)) \rangle$;
- (c) $\langle \xi \otimes \eta \otimes \lambda, [\text{Ad}(g_1(\lambda))(t^{13}), [(d_3 g_2) g_2^{-1}, \text{Ad}(g_2(\lambda))(\lambda_2^{\vee})]] \rangle = \langle \text{ad}^*(\bar{x})^{\otimes 2}(\xi \otimes \eta), \text{Ad}(g)^{\otimes 2}(g_2^{-1} d_1(g_2)) \rangle$;
- (d)

$$\begin{aligned} & \langle \xi \otimes \eta \otimes \lambda, [(d_3 g_1) g_1^{-1}, \text{Ad}(g_1(\lambda))(\lambda_1^{\vee})], [(d_3 g_2) g_2^{-1}, \text{Ad}(g_2(\lambda))(\lambda_2^{\vee})] \rangle \\ &= \langle \text{ad}^*(\bar{x})(\xi) \otimes \text{ad}^*(\bar{x})(\eta) \otimes \lambda, [(d_3 g_1) g_1^{-1}, (d_3 g_2) g_2^{-1}] \rangle. \end{aligned}$$

We have $\{F_{\xi}, F_{\eta}\}'_G(g) = \{f_{\xi}, f_{\eta}\}_{\mathfrak{g}^*}(a^{-1}(g))$; then $\bar{x} = x$, so $\{F_{\xi}, F_{\eta}\}'_G = \{F_{\xi}, F_{\eta}\}_G$ iff

$$\begin{aligned} & \langle \text{ad}^*(x)(\xi) \otimes \text{ad}^*(x)(\eta), \text{Ad}(g)^{\otimes 2}(g_2^{-1} d_1(g_2) - g_1^{-1} d_2(g_1)) \rangle \\ &+ \langle \text{id} \otimes \text{id} \otimes \lambda, [d_3(g_1) g_1^{-1}, d_3(g_2) g_2^{-1}] \rangle - (\text{id} \otimes \varphi(\text{ad}(x)))(t) - r_0 = 0, \end{aligned}$$

for which a sufficient condition is that $g(\lambda)$ satisfies (2).

Remark 1.3. Formula (4) implies that the image of $\{-, -\}_G$ under the map $\log : G \rightarrow \mathfrak{g}$ is given by

$$\{f, g\}(x) = \langle df(x) \otimes dg(x), (\text{ad}(x) \otimes (\frac{1}{2} \text{ad}(x) \coth(\frac{1}{2} \text{ad}(x))))(t) + \text{ad}(x)^{\otimes 2}(r_0) \rangle;$$

this is a result of [FM2].

1.2. Construction based on quantization of Lie bialgebras. We now give a proof of Theorem 0.1 based on the theory of quantization of Lie bialgebras.

1.2.1. Construction of $g(\lambda)$ (proof of Proposition 0.2). Set $U := U(\mathfrak{g})[[\hbar]]$, $U' := \{x \in U \mid \forall n \geq 0, \delta_n(x) \in \hbar^n U^{\widehat{\otimes} n}\} = U(\hbar \mathfrak{g}[[\hbar]]) \subset U$ (here $\delta_n = (\text{id} - \eta \circ \varepsilon)^{\otimes n} \circ \Delta^{(n)}$, η is the unit map $\mathbf{k}[[\hbar]] \rightarrow U$). As a $\mathbf{k}[[\hbar]]$ -algebra, U' is a flat deformation of $\widehat{S}(\mathfrak{g}) = \mathbf{k}[[\mathfrak{g}^*]]$.

Let $\Phi \in U^{\widehat{\otimes} 3}$ be an admissible associator. This means that

$$\Phi \in 1 + \frac{\hbar^2}{24} [t^{1,2}, t^{2,3}] + \hbar^2 U^{\widehat{\otimes} 3}, \quad \hbar \log(\Phi) \in (U')^{\widehat{\otimes} 3},$$

$(U, m, \Delta_0, \mathcal{R}_0 = 1, \Phi)$ is a quasitriangular quasi-Hopf algebra. We also require $\varepsilon^{(i)}(\Phi) = 1$, $i = 1, 2, 3$ (here $\varepsilon^{(1)} = \varepsilon \otimes \text{id} \otimes \text{id}$, etc., ε is the counit, m and Δ_0 are the undeformed product and coproduct of U). According to [EH], any universal Lie associator gives rise to an admissible associator.

According to [EK], there exists a twist killing Φ , and according to [EH], this twist can then be made admissible by a suitable gauge transformation. The resulting twist J satisfies the following conditions: $J \in U^{\widehat{\otimes} 2}$, $J = 1 - \hbar r/2 + o(\hbar)$, $\hbar \log(J) \in (U')^{\widehat{\otimes} 2}$, $(\varepsilon \otimes \text{id})(J) = (\text{id} \otimes \varepsilon)(J) = 1$,

$$\Phi = (J^{2,3} J^{1,23})^{-1} J^{1,2} J^{12,3}. \quad (5)$$

Then $U^J := (U, m, \Delta^J, \mathcal{R})$ is a quasitriangular Hopf algebra quantizing $(\mathfrak{g}, r)$. Here $\Delta^J(x) = J \Delta_0(x) J^{-1}$ and $\mathcal{R} = J^{2,1} e^{\hbar t/2} J^{-1}$.

We have $\text{Ker}(\varepsilon) \cap U' \subset \hbar U$, therefore $\log(J) \in U \widehat{\otimes} U'$. Then its reduction mod $\hbar$, denoted $g(\lambda) = g^{1,2} = \log(J)|_{\hbar=0}$, belongs to $U(\mathfrak{g}) \widehat{\otimes} \widehat{S}(\mathfrak{g}) = U(\mathfrak{g})[[\mathfrak{g}^*]]$ (formal series on $\mathfrak{g}^*$ with coefficients in $U(\mathfrak{g})$). The reduction mod $\hbar$ of (5) is $g^{12,3} = g^{1,3} g^{2,3}$. Since we also have $(\varepsilon \otimes \text{id})(g) = 1$, we get $g = \exp(A)$, with $A \in \mathfrak{g} \otimes \widehat{S}(\mathfrak{g})_{>0}$.

Lemma 1.4. *$g(\lambda)$ satisfies (2).*

Proof. According to [EE], $\Phi \in U^{\widehat{\otimes} 2} \widehat{\otimes} U'$ has the expansion $1 + \hbar \phi_1 + o(\hbar^2)$, where $\phi_1 \in U^{\widehat{\otimes} 2} \widehat{\otimes} U'$ is such that $(\phi_1 - \phi_1^{2,1,3})|_{\hbar=0} = -\rho_{\text{AM}}$.

If $x \in U(\mathfrak{g}) \widehat{\otimes} \widehat{S}(\mathfrak{g})$, we denote by $\bar{x}$ a lift of x in $U \widehat{\otimes} U'$.

Let us expand $\log(J)$ as $\bar{A} + \hbar A_1 + o(\hbar)$, with $A_1 \in U \widehat{\otimes} U'$.

Then $J^{1,23} = \exp(\bar{A}^{1,3} + \hbar(A_1^{1,3} + d_2 A^{1,3}) + o(\hbar)) = J^{1,3}(1 + \hbar \overline{g_1^{-1} d_2(g_1)} + o(\hbar))$.

We have $[J^{1,3}, J^{2,3}] = \hbar \{g^{1,3}, g^{2,3}\} + o(\hbar)$, so $(J^{12,3})^{-1} [J^{1,3}, J^{2,3}] = \hbar \overline{(g^{1,3} g^{2,3})^{-1} \{g^{1,3}, g^{2,3}\}} + o(\hbar) = \hbar \langle \text{id} \otimes \text{id} \otimes \lambda, [g_1^{-1} d_3(g_1), g_2^{-1} d_3(g_2)] \rangle + o(\hbar)$. So we get

$$J^{12,3} = J^{2,3} J^{1,3} (1 + \hbar \psi_1 + o(\hbar)),$$

where $\psi_1 \in U^{\widehat{\otimes} 2} \widehat{\otimes} U'$ is such that $(\psi_1 - \psi_1^{2,1,3})|_{\hbar=0} = \langle \text{id} \otimes \text{id} \otimes \lambda, [g_1^{-1} d_3(g_1), g_2^{-1} d_3(g_2)] \rangle$.

Then (5) gives

$$1 + \hbar \phi_1 + o(\hbar) = (1 - \hbar \overline{g_1^{-1} d_2(g_1)} + o(\hbar)) (J^{1,3})^{-1} (J^{2,3})^{-1} (1 - \hbar r/2 + o(\hbar)) J^{2,3} J^{1,3} (1 + \hbar \psi_1 + o(\hbar)).$$

The reduction modulo $\hbar$ of $(J^{1,3})^{-1} (J^{2,3})^{-1} r J^{2,3} J^{1,3}$ is $\text{Ad}(g \otimes g)^{-1}(r)$. Then subtracting 1, dividing by $\hbar$, reducing modulo $\hbar$ and antisymmetrizing the two first tensor factors, we get the lemma. $\square$

More generally, assume that $J' \in U^{\widehat{\otimes} 2}$ satisfies $J' = 1 - \hbar r/2 + o(\hbar)$, $\hbar \log(J') \in (U')^{\widehat{\otimes} 2}$, $(\varepsilon \otimes \text{id})(J') = 1$ and

$$\Phi = (J'^{2,3} J'^{1,23})^{-1} J'^{1,2} J'^{12,3}. \quad (6)$$

Then $J' \in U \widehat{\otimes} U'$, and its reduction g' modulo $\hbar$ satisfies: $g' \in \exp(\mathfrak{g} \otimes \widehat{S}(\mathfrak{g})_{>0})$, and equation (2) with g replaced by g' . Moreover, there exists $u \in U^\times$ such that $J' = u^2 J(u^{12})^{-1}$.

Remark 1.5. Equation (6) can be interpreted as saying that J' is a vertex-IRF transformation relating $J^{1,2}$ and Φ , and equation (2) for g' is the classical limit of this statement (see [EN]). Vertex-IRF transformations are a special kind of non-invariant dynamical gauge transformations, which map a constant, but non-invariant twist to an invariant, but non-constant (i.e., dynamical) one. $\square$

Let $U'_0 := \text{Ker}(\varepsilon) \cap U'$. Then $\hbar^{-1} U'_0 \subset U$ is a Lie subalgebra for the commutator. This Lie algebra acts on the set of solutions of (6) by $\delta_u(J) = u^2 J - J u^{12}$ ($\delta_u(J) \in U \widehat{\otimes} U'$ because it is equal to $[u^2, J] - J(u^{12} - u^2)$); this means that if ε is a formal parameter with $\varepsilon^2 = 0$, then $(\text{id} + \varepsilon \delta_u)(J')$ is a solution of (6) if J' is.

The reduction modulo $\hbar$ of this action may be described as follows. The Lie algebra $(\widehat{S}(\mathfrak{g})_{>0}, \{-, -\})$ acts on the set of solutions of (2) by $\delta_f(g) = \{1 \otimes f, g\} - g \cdot df$, i.e., action (3).

When restricted to the Lie subalgebra $\widehat{S}(\mathfrak{g})_{>1}$, this action is the infinitesimal of the right action of $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ on the set of solutions of (2), given by $(g * \alpha)(\lambda) = g(\text{Ad}^*(\alpha(\lambda))(\lambda))\alpha(\lambda)$.

1.2.2. Isomorphism $\mathfrak{g}^* \rightarrow G^*$ (proof of Proposition 0.3). It follows from [EH] that the subalgebras $(U^J)'$ and U' of U are equal. According to [Dr1, Gav], $(U^J)'$ is a flat deformation of $\mathcal{O}_{G^*} := U(\mathfrak{g}^*)^*$. On the other hand, U' is a flat deformation of $\mathcal{O}_{\mathfrak{g}^*} := \widehat{S}(\mathfrak{g})$. So the equality $i_{\hbar} : (U^J)' \rightarrow U'$ induces an isomorphism of Poisson algebras $i : \mathcal{O}_{G^*} \xrightarrow{\sim} \mathcal{O}_{\mathfrak{g}^*}$, and therefore a Poisson isomorphism $\mathfrak{g}^* \rightarrow G^*$.

To express this isomorphism explicitly, we will construct maps $a : \mathcal{O}_G \rightarrow \mathcal{O}_{\mathfrak{g}^*}$ and $b : \mathcal{O}_G \rightarrow \mathcal{O}_{G^*}$, such that $i = a \circ b^{-1}$.

We first construct quantized versions of a and b . We have $\hbar \log(J) \in (U')^{\otimes 2}$, therefore $J, J^{-1} \in U \widehat{\otimes} U'$ and $Je^{\hbar t} J^{-1} \in U \widehat{\otimes} U'$. We then define $a_{\hbar} : U^* \rightarrow U'$ by $a_{\hbar}(f) := (f \otimes \text{id})(Je^{\hbar t} J^{-1})$. Now we also have $J^{2,1} \in U \widehat{\otimes} U'$, hence $\mathcal{R}, \mathcal{R}^{2,1} \in U \widehat{\otimes} U' = U \widehat{\otimes} (U^J)'$. We then define the linear map $b_{\hbar} : U^* \rightarrow (U^J)'$ by $b_{\hbar}(f) := (f \otimes \text{id})(\mathcal{R}^{2,1} \mathcal{R})$. Then $a_{\hbar} \circ b_{\hbar}^{-1} : U' \rightarrow (U^J)'$ coincides with $i_{\hbar}$. We define a, b as the reductions modulo $\hbar$ of $a_{\hbar}, b_{\hbar}$.

Let us now compute the classical limit of $a_{\hbar}$. Define maps $j_{\hbar}, \iota_{\hbar}, j'_{\hbar} : U^* \rightarrow U'$ by $j_{\hbar}(f) = (f \otimes \text{id})(J)$, $j'_{\hbar}(f) = (f \otimes \text{id})(J^{-1})$. Recall that the reductions modulo $\hbar$ of all three elements $J, J^{-1}, e^{\hbar t}$ in $U(\mathfrak{g}) \widehat{\otimes} \widehat{S}(\mathfrak{g})$ are of the form $K = \exp(k)$, where $k \in \mathfrak{g} \otimes \widehat{S}(\mathfrak{g})_{>0}$.

Lemma 1.6. *If $K \in U(\mathfrak{g}) \widehat{\otimes} \widehat{S}(\mathfrak{g})$ is of the form $\exp(k)$, where $k \in \mathfrak{g} \otimes \widehat{S}(\mathfrak{g})_{>0}$, then the morphism $\mathcal{O}_G = U(\mathfrak{g})^* \rightarrow \widehat{S}(\mathfrak{g}) = \mathcal{O}_{\mathfrak{g}^*}$ given by $f \mapsto (f \otimes \text{id})(K)$ is dual to the morphism $\mathfrak{g}^* \rightarrow G$, $\lambda \mapsto e^{k(\lambda)}$.*

Proof. We compose this morphism with the transpose of the inverse of the symmetrization map ${}^t \text{Sym}^{-1} : S(\mathfrak{g})^* \rightarrow U(\mathfrak{g})^*$. The morphism ${}^t \text{Sym}^{-1}$ corresponds to the logarithm map $G \rightarrow \mathfrak{g}$. Now the composed morphism $S(\mathfrak{g})^* \rightarrow \widehat{S}(\mathfrak{g})$ is given by $f \mapsto (f \otimes \text{id})((\text{Sym}^{-1} \otimes \text{id})(K))$. Now $(\text{Sym}^{-1} \otimes \text{id})(K)$ is $\exp(k)$, where the exponential is now taken in $S(\mathfrak{g}) \widehat{\otimes} \widehat{S}(\mathfrak{g})$. The morphism $S(\mathfrak{g})^* \rightarrow \widehat{S}(\mathfrak{g})$, $f \mapsto (f \otimes \text{id})(\exp(k))$, is an algebra morphism, taking the function $X \mapsto \alpha(X)$ on $\mathfrak{g}$ ($\alpha \in \mathfrak{g}^*$) to the function $\lambda \mapsto \alpha(k(\lambda))$ on $\mathfrak{g}^*$, and therefore corresponds to the morphism $\mathfrak{g}^* \rightarrow \mathfrak{g}$, $\lambda \mapsto k(\lambda)$. Composing with the exponential, we get the announced morphism. $\square$

It follows that the reductions modulo $\hbar$ of the morphisms $j_{\hbar}, \iota_{\hbar}, j'_{\hbar}$ are morphisms $\mathcal{O}_G \rightarrow \mathcal{O}_{\mathfrak{g}^*}$, corresponding to morphisms $j, \iota, j' : \mathfrak{g}^* \rightarrow G$ such that $j(\lambda) = g(\lambda)$, $\iota(\lambda) = e^{\lambda^\vee}$, $j'(\lambda) = g(e^{\lambda})^{-1}$.

Then $a_{\hbar} = m^{(2)} \circ (j_{\hbar} \otimes \iota_{\hbar} \otimes j'_{\hbar}) \circ \Delta^{(2)}$, so $a : \mathcal{O}_{G^*} \rightarrow \mathcal{O}_G$ corresponds to the composed map $G^* \xrightarrow{\text{diag}^{(2)}} (G^*)^3 \xrightarrow{(j, \iota, j')} G^3 \xrightarrow{\text{product}^{(2)}} G$. Therefore we get $a(\lambda) = g(\lambda) e^{\lambda^\vee} g(\lambda)^{-1}$.

We now compute the classical limit of $b_{\hbar}$. Define maps $L_{\hbar}, R'_{\hbar} : (U^J)^* \rightarrow (U^J)'$ by $L_{\hbar}(f) := (f \otimes \text{id})(\mathcal{R}^{2,1})$, $R'_{\hbar}(f) := (f \otimes \text{id})(\mathcal{R})$. Then $L_{\hbar}$ is an antimorphism of algebras and a morphism of coalgebras, while $R'_{\hbar}$ is a morphism of algebras and antimorphism of coalgebras. Their reductions L, R' modulo $\hbar$ are morphisms $\mathcal{O}_{G^*} \rightarrow \mathcal{O}_G$ (anti-Poisson coalgebra morphism in the case of L , Poisson anti-coalgebra in the case of R'). Using [EGH], appendix, one shows that these morphisms correspond to the morphisms of formal groups $L, R' : G^* \rightarrow G$ (antimorphism in the case of R'), corresponding to the morphisms $L, R' : \mathfrak{g}^* \rightarrow \mathfrak{g}$, given by $L(\lambda) := (\lambda \otimes \text{id})(r)$ and $R'(\lambda) := (\lambda \otimes \text{id})(r^{2,1})$. Here L is a Lie algebra, anti-Lie coalgebra morphism and R' is an anti-Lie algebra, Lie coalgebra morphism. Now $b_{\hbar} = m \circ (L_{\hbar} \otimes R_{\hbar}) \circ \Delta$, so $b : \mathcal{O}_{G^*} \rightarrow \mathcal{O}_G$ corresponds to the composed map $G^* \xrightarrow{\text{diag}} (G^*)^2 \xrightarrow{(L, R')} G^2 \xrightarrow{\text{product}} G$, i.e., $g^* \mapsto L(g^*) R'(g^*) = L(g^*) R(g^*)^{-1}$.

Finally, the isomorphism $i : \mathfrak{g}^* \rightarrow G^*$ is equal to $b^{-1} \circ a$, so it takes $\lambda \in \mathfrak{g}^*$ to $g^* \in G^*$ such that

$$L(g^*) R(g^*)^{-1} = g(\lambda) e^{\lambda^\vee} g(\lambda)^{-1}.$$

So it coincides with the isomorphism obtained in Proposition 0.3.

1.3. On the group $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$. In this section, we give a "quantum" proof of the following statement, which was used in Section 1.1 (and can also be proved in the setup of this section).

Proposition 1.7. $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ is a subgroup of $\text{Map}_0(\mathfrak{g}^*, G)$.

Proof. Set $U'_0 = \text{Ker}(\varepsilon) \cap U'$. The map of $U' \rightarrow \widehat{S}(\mathfrak{g})$ of reduction by $\hbar$ restricts to $U'_0 \rightarrow \widehat{S}(\mathfrak{g})_{>0}$, so we get a map $U'_0 \rightarrow \mathfrak{g}$.

Consider the set of all J' , such that $\hbar \log(J') \in (U'_0)^{\otimes 2}$, the image of $\hbar \log(J')$ in $\mathfrak{g}^{\otimes 2}$ is zero, and $J'^{2,3} J'^{1,23} = J'^{1,23}$.

This is the set of elements of the form $J'(u) = u^2(u^{12})^{-1}$, where $u \in \exp(U'_0/\hbar)$ is such that the image of $\log(u)$ under $\widehat{S}(\mathfrak{g})_{>0} \rightarrow \mathfrak{g}$ is zero. Indeed, one recovers u from J' by $u = (\text{id} \otimes \varepsilon)(J')^{-1}$.

We denote by $A(\lambda)$ the image of $\hbar \log(J')$ by $\widehat{S}(\mathfrak{g})_{>0} \widehat{\otimes} \widehat{S}(\mathfrak{g})_{\geq 2} \rightarrow \mathfrak{g} \otimes \widehat{S}(\mathfrak{g})_{\geq 2}$, and set $g(\lambda) := \exp(A(\lambda))$. Then $g(\lambda) \in \text{Map}_0(\mathfrak{g}^*, G)$.

As in Lemma 1.4, one proves that $g(\lambda)$ satisfies equation (1). Hence we get a map of sets $\{J' \text{ as above}\} \rightarrow \text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$, which is surjective.

On the other hand, the set of all $J'(u)$ is equipped with a product, such that $J'(u) * J'(v) = J'(uv)$. This product expresses as follows: $J'(u) * J'(v) = (u^2 J'(v)(u^2)^{-1}) J'(u)$; the classical limit of this expression is the product formula for $\text{Map}_0(\mathfrak{g}^*, G)$. So $\text{Map}_0^{\text{ham}}(\mathfrak{g}^*, G)$ is a subgroup of $\text{Map}_0(\mathfrak{g}^*, G)$. $\square$

2. QUANTIZATION OF ρ_{FM}

In this section, we prove Theorem 0.4.

Let Φ_{univ} be a universal Lie associator defined over $\mathbf{k}$ with parameter $\mu = 1$. So $\Phi_{\text{univ}} = 1 + \frac{1}{24}[t_{12}, t_{23}] + \text{terms of degree } > 2$, where t_{ij} is the universal version of $t^{i,j}$ (see [Dr3]). Set $\Phi_\nu := \Phi_{\text{univ}}(2\hbar\nu t^{1,2}, 2\hbar\nu t^{2,3})$ and $\Phi := \Phi_{1/2}$. Then

$$(U(\mathfrak{g})[[\hbar]], m, \Delta_0, \Phi_\nu)$$

is a quasi-Hopf algebra; its classical limit is the quasi-Lie bialgebra $(\mathfrak{g}, \delta = 0, \nu^2[t^{1,2}, t^{2,3}])$. Let J be an admissible twist killing Φ , and let us twist this quasi-Hopf algebra by J . We obtain the quasi-Hopf algebra

$$(U(\mathfrak{g})[[\hbar]], m, \Delta_0^J, \Phi_\nu^J),$$

where $\Delta_0^J(x) = J\Delta_0(x)J^{-1}$ and $\Phi_\nu^J = J^{2,3}J^{1,23}\Phi_\nu(J^{-1})^{12,3}(J^{-1})^{1,2}$. Its classical limit is the quasi-Lie bialgebra $(\mathfrak{g}, \delta(x) = [x^1 + x^2, r], Z_\nu)$.

Now $(U(\mathfrak{g})[[\hbar]], m, \Delta_0^J)$ is a Hopf algebra quantizing $(\mathfrak{g}, \delta)$, which we denote by $U_\hbar(\mathfrak{g})$, and Φ_ν^J is an associator for this quantized universal enveloping algebra, with classical limit Z_ν .

Φ_ν^J clearly satisfies the invariance and pentagon equations. We have $J = 1 + \hbar j_1$, where $j_1 \in U \widehat{\otimes} U'$, so $J^{2,3}, J^{1,23}, (J^{-1})^{12,3}$ and $(J^{-1})^{1,2}$ are all of the form $1 + \hbar k$, $k \in U^{\widehat{\otimes} 2} \widehat{\otimes} U'$. Hence $\Phi_\nu^J = 1 + \hbar \psi_\nu$, where $\psi_\nu \in U^{\widehat{\otimes} 2} \widehat{\otimes} U'$. Set $U_\hbar := U_\hbar(\mathfrak{g})$, then $U'_\hbar = U'$, so $\psi_\nu \in U_\hbar^{\widehat{\otimes} 2} \widehat{\otimes} U'_\hbar$. We now compute $(\psi_\nu - \psi_\nu^{2,1,3})|_{\hbar=0}$; this is an element of $U(\mathfrak{g})^{\otimes 2} \widehat{\otimes} \mathcal{O}_{G^*} \simeq U(\mathfrak{g})^{\otimes 2} \widehat{\otimes} \mathcal{O}_{\mathfrak{g}^*}$.

We have $\Phi_\nu^J = \Phi^J + (\Phi_\nu - \Phi)^J = 1 + \hbar(\Phi_\nu - \Phi)^J$. Let us define ϕ, ϕ_ν by $\Phi = 1 + \hbar\phi$, $\Phi_\nu = 1 + \hbar\phi_\nu$, then $(\psi_\nu)|_{\hbar=0} = ((\phi_\nu - \phi)^J)|_{\hbar=0}$, therefore

$$\begin{aligned} (\psi_\nu - \psi_\nu^{2,1,3})|_{\hbar=0} &= ((\phi_\nu - \phi_\nu^{2,1,3})^J)|_{\hbar=0} - ((\phi - \phi^{2,1,3})^J)|_{\hbar=0} \\ &= \text{Ad}(g(\lambda))^{\otimes 2}(\rho_{\text{AM}}(\lambda) - \rho_{\text{AM}}^\nu(\lambda)) \end{aligned}$$

as a formal function $\mathfrak{g}^* \rightarrow \wedge^2(\mathfrak{g})$. Here $\rho_{\text{AM}}^\nu(\lambda) = 2\nu\rho_{\text{AM}}(2\nu\lambda)$. Since ρ_{AM} and ρ_{AM}^ν are G -equivariant, this is

$$(\rho_{\text{AM}} - \rho_{\text{AM}}^\nu)(\text{Ad}^*(g(\lambda))(\lambda)) = \left(\text{id} \otimes \left(\frac{1}{2} \frac{e^{\text{ad}(\bar{\lambda}^\vee)} + \text{id}}{e^{\text{ad}(\bar{\lambda}^\vee)} - \text{id}} - \nu \frac{e^{2\nu \text{ad}(\bar{\lambda}^\vee)} + \text{id}}{e^{2\nu \text{ad}(\bar{\lambda}^\vee)} - \text{id}} \right) \right)(t) = \rho_{\text{FM}}(g^*).$$

Here we set $\bar{\lambda} := \text{Ad}^*(g(\lambda))(\lambda)$ and we use the relation $L(g^*)G(g^*)^{-1} = e^{\bar{\lambda}^\vee}$.

We have proved:

Theorem 2.1. *The quantized universal enveloping algebra $U_\hbar(\mathfrak{g}) = U(\mathfrak{g})[[\hbar]]^J$, together with the pair $(\bar{J}, \bar{\Phi})$ defined by $\bar{J} = \bar{\Phi} = \Phi_\nu^J$, is a quantization of the Poisson-Lie dynamical r -matrix $\rho_{\text{FM}}(g^*)$ over $(G^*, \mathfrak{g}, Z_\nu)$.*

REFERENCES

- [Al] A. Alekseev, *On Poisson actions of compact Lie groups on symplectic manifolds*, J. Differential Geom. **45** (1997), 241-56.
- [AM] A. Alekseev, E. Meinrenken, *The non-commutative Weil algebra*, Invent. Math. **139** (2000), 135-72.
- [BDF] J. Balog, L. Dabrowski, and L. Fehér, *Classical r -matrix and exchange algebra in WZNW and Toda field theories*, Phys. Lett. B, **244** (1990), issue 2, 227-34.
- [BFP] J. Balog, L. Fehér, L. Palla, *Chiral extensions of the WZNW phase space, Poisson-Lie symmetries and groupoids*, preprint hep-th/9910046, Nucl. Phys. B **568** (2000), 503-42.
- [Bo] P. Boalch, *Stokes matrices, Poisson-Lie groups and Frobenius manifolds*, Invent. Math. **146** (2001), no. 3, 479-506.
- [Ch] V. Chloup-Arnould, *Linearization of some Poisson-Lie tensor*, J. Geom. Phys. **24** (1997), no. 1, 46-52.
- [Dr1] V. Drinfeld, *Quantum groups*, Proceedings of the ICM-86 (Berkeley), 798-820, Amer. Math. Soc., Providence, RI, 1987.
- [Dr2] V. Drinfeld, *Quasi-Hopf algebras*, Leningrad Math. J. **1** (1990), no. 6, 1419-57.
- [Dr3] V. Drinfeld, *On quasitriangular quasi-Hopf algebras and on a group that is closely connected with $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$* , Leningrad Math. J. **2** (1991), no. 4, 829-60.
- [EE] B. Enriquez, P. Etingof, *Quantization of Alekseev-Meinrenken dynamical r -matrices*, preprint math.QA/0302067 (in memory of F.I. Karpelevich), AMS Transl. **210** (2003), no. 2, 81-98.
- [EEM] B. Enriquez, P. Etingof, I. Marshall, *Quantization of some Poisson-Lie dynamical r -matrices and Poisson homogeneous spaces*, preprint math.QA/0403283 (in memory of J. Donin).
- [EGH] B. Enriquez, F. Gavarini, G. Halbout, *Uniqueness of braidings of quasitriangular Lie bialgebras and lifts of classical r -matrices*, Internat. Math. Res. Notices, **46** (2003), 2461-86.
- [EH] B. Enriquez, G. Halbout, *Poisson algebras associated to quasi-Hopf algebras*, Adv. Math. **186** (2004), no. 2, 363-95.
- [EK] P. Etingof, D. Kazhdan, *Quantization of Lie bialgebras, I, II*, Selecta Math. (N.S.) **2** (1996), no. 1, 1-41; **4** (1998), no. 2, 213-31.
- [EN] P. Etingof, D. Nikshych, *Vertex-IRF transformations and quantization of dynamical r -matrices*, Math. Res. Lett. **8** (2001), no. 3, 331-45.
- [ES] P. Etingof, O. Schiffmann, *On the moduli space of classical dynamical r -matrices*, Math. Res. Lett. **8** (2001), no. 1-2, 157-70.
- [FM1] L. Fehér, I. Marshall, *On a Poisson-Lie analogue of the classical dynamical Yang-Baxter equation for self-dual Lie algebras*, Lett. Math. Phys. **62** (2002), 51-62.
- [FM2] L. Fehér, I. Marshall, *The non-abelian momentum map for Poisson-Lie symmetries on the chiral WZNW phase space*, Int. Math. Res. Not. **49** (2004), 2611-36.
- [Gav] F. Gavarini, *The quantum duality principle*, Ann. Inst. Fourier (Grenoble) **52** (2002), no. 3, 809-34.
- [GW] Viktor L. Ginzburg, A. Weinstein, *Lie-Poisson structure on some Poisson-Lie groups*, J. Amer. Math. Soc. **5:2** (1992), 445-53.
- [STS] M. Semenov-Tian-Shansky, *Dressing transformations and Poisson group actions*, Publ. Res. Inst. Math. Sci. **21** (1985), no. 6, 1237-1260.

IRMA (CNRS), 7 RUE RENÉ DESCARTES, F-67084 STRASBOURG, FRANCE

E-mail address: `enriquez@math.u-strasbg.fr`

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

E-mail address: `etingof@math.mit.edu`

DÉPARTEMENT DE MATHÉMATIQUES, EPFL, CH-1015 LAUSANNE, SWITZERLAND

E-mail address: `ian.marshall@epfl.ch`